**Open Access** 

# Machine learning prediction model of prolonged delay to loop ileostomy closure after rectal cancer surgery: a retrospective study

Jianying Liu<sup>1</sup>, Mengxiao Jiang<sup>2</sup>, Xiaoping Chen<sup>2</sup>, Yonglan Ge<sup>1</sup>, Zongxin Zheng<sup>2</sup>, Xia Yang<sup>1</sup>, Wenhao Zhou<sup>1</sup>, Huiting Zhang<sup>3\*</sup>, Meichun Zheng<sup>1\*</sup> and Baojia Luo<sup>1\*</sup>

# Abstract

**Background** Delayed closure of a temporary ileostomy in patients with rectal cancer may cause psychological, physiological, and socioeconomic burdens to patients.

**Purpose** This study aimed to develop and validate a machine learning-based model to predict the delayed ileostomy closure after surgery in patients with rectal cancer.

Design A retrospective study.

**Methods** LASSO regression was used for feature screening, and XGBoost was used for machine learning model construction. Model performance was assessed by receiver operating characteristic (ROC) curve analysis, calibration curve analysis, clinical decision curve analysis, sensitivity, specificity, accuracy, and F1 score. The SHAP method was used to interpretate the results of the machine learning model.

**Results** A total of 442 rectal cancer patients who received a loop ileostomy were included in this study, and 305 experienced delayed closure (69%). The XGBoost model area under the ROC curve (AUC) of the training set was 0.744 (95% confidence interval [CI]: 0.686–0.806) and of the test set was 0.809 (95% CI: 0.728–0.889). The importance of each variable, in descending order was body mass index (BMI), postoperative chemotherapy, distance from tumor to anal margin, depth of tumor infiltration, neoadjuvant chemoradiotherapy, and anastomotic stenosis. The importance of SHAP variables in the model from high to low was: 'BMI'' postoperative chemotherapy'' distance of the tumor from the anal verge'' depth of tumor infiltration'' neoadjuvant radiotherapy'' anastomotic stenosis'.

<sup>†</sup>Jianying Liu and Mengxiao Jiang contributed equally to this work.

\*Correspondence: Huiting Zhang zhanght@sysucc.org.cn Meichun Zheng zhengmch@sysucc.org.cn Baojia Luo Iuobj@sysucc.org.cn

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit ine to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by-nc-nd/4.0/.



**Conclusion** The XGBoost machine learning model we constructed showed good performance in predicting delayed closure of loop ileostomy in rectal cancer patients. In addition, the SHAP method can help better understand the results of machine learning models.

Keywords Machine learning, Rectal cancer, Loop ileostomy, Delayed closure

# Background

Rectal cancer (RC) is one of the most common malignant digestive tract tumors. colorectal cancer (CRC) ranks third globally in incidence and is the second leading cause of cancer-related death [1]. Total mesorectal excision (TME), proposed by Heald et al. in 1982 [2], remains the standard for radical RC resection. Its basic principle is to completely remove the mesentery surrounding the tumor at the anatomical and embryonic levels. Over time, techniques have advanced, with laparoscopic transabdominal tumor excision (laTME) becoming widely used for resectable cases [3]. After rectal cancer surgery, patients may experience postoperative complications, including anastomotic leakage, which can influence recovery, increase healthcare utilization, and potentially affect oncological outcomes [4–5]. To reduce the risk of clinical consequences association with leakage, a diverting ileostomy is often performed at the time of intestinal anastomoses [6].

However, the effectiveness of a temporary ileostomy in preventing or reduceing anastomotic leakage remains debated. Many studies support its role in preventing anastomotic leakage [7-8]. Garg et al. [9] conducted a meta-analysis of 5 randomized controlled trials including 768 patients, showing that ileostomy significantly reduced the leakage and reoperation rates. In recent decades, prophylactic stoma has been widely used in anal sphincter preservation surgery, especially for low rectal cancer [10]. Conversely, other studies suggest that ileostomy may not reduce leakage but can reduce the risk of abdominal infection [11-13]. Currently, no standardized guideline exist for whether patients should undergo temporary ileostomy. To reduce postoperative complications, the use of temporary preventive ileostomies has increased, leading to unnecessary stoma related complications in patients without anastomotic leakage.

Temporary ileostomy can negatively affect patients' psychological well-being and is associated with various stoma-related complications [14]. These include intestinal obstruction, diarrhea, anastomotic leakage, incision infection, and enterocutaneous fistulae [15]. An international study involving 279 ileostomy patients reported an overall complication rate of 83% [16]. As such, a temporary ileostomy should be closed as soon as possible. Delayed closure beyond 6 months is associated with a 3.7-fold increased risk of severe intestinal dysfunction after restoring intestinal continuity [17]. In addition, the incidence of stoma-related complications increases with prolonged closure intervals [18]. Patients who undergo

ileostomy closure more than 3 months after rectal surgery experience a significant decline in their quality of life [19–20] and higher healthcare cost [21]. Given the variability in closure timing, a predictive model to identify patients at risk of delayed closure would aid clinical decision making.

Machine learning (ML) is increasingly used in healthcare to assess patient risk and support decision making [22–24]. SHapley Additive exPlans (SHAP) is an advanced, interpretable ML framework designed to provide in-depth explanations for the predictions of any ML model [25–26]. Other stud using broader populationbased datasets have found it more challenging to achieve accurate prediction [27]. ML is currently widely used in the diagnosis, treatment, and prediction of cancer recurrence, and models have practical applications [28–30]. Therefore, this study aimed to develop an interpretable ML model using SHAP to predict the risk of delayed closure of an ileostomy after rectal cancer surgery.

# Methods

# Study setting and patients

An observational prospective cohort study was conducted in a large oncology hospital in China. The data of patients who underwent TME for rectal cancer and received a temporary ileostomy at the Colorectal Surgery Department of Sun Yat-sen University Cancer Center from January 1, 2022 to December 31, 2023 were retrospectively collected and reviewed. This study was approved by the Ethics Review Committee of Sun Yatsen University Cancer Center. This study has obtained informed consent from the participants and signed the informed consent form.

# Inclusion criteria and exclusion criteria

Inclusion criteria were: (1) Diagnosed with rectal cancer by colonoscopy and pathological examination of a biopsy tissue specimen; (2) Received an ileostomy; (3) Regular follow-up at Sun Yat-sen University Cancer Center; (4) No tumor recurrence during follow-up. Exclusion criteria were: (1) Diabetes mellitus or hypertension; (2) Ongoing steroid treatment; (3) Immune or inflammatory diseases such as inflammatory bowel disease and autoimmune diseases; (4) Concomitant small intestine or colon resection during surgery; (5) Repeat or multiple surgeries during the period from rectal cancer surgery to ileostomy closure.

## Feature collection and screening

Prior to data collection, we conducted a comprehensive literature review to identify potential factors that may delay ileostomy closure. Patient data were collected from the standardized database of Sun Yat-sen University Cancer Center, and included age, sex, body mass index (BMI), distance between tumor and anal margin, pathological type, depth of tumor infiltration, lymph node metastasis, distant metastases, preoperative neoadjuvant chemoradiotherapy, postoperative chemotherapy, anastomotic leakage, anastomotic stenosis, low anterior resection syndrome (LARS), complications related to ileostomy. Variables were selected by LASSO regression (n-fold = 20) using the "glmnet" package in R.

## Model development and validation

The data set was partitioned before building the model: 75% of the patients were assigned to the training set using the "caret" package, and the remaining 25% were assigned to the validation set. The model was built using the "XGBoost" package, and used the train() function to optimize the parameters in the caret package, and output the optimal parameter configuration. The XGBoost model is constructed by setting the learning rate eta to 0.1, the maximum depth (max\_depth) to 2, and the number of iteration rounds (i.e., enhancement rounds) to 100. Accuracy, sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and F1 score, ROC curve, calibration curve, and clinical decision curve analysis (DCA) were used to evaluate the performance of the model in the training and test sets. Predicted outcome probabilities were converted to binary outcomes using a threshold of 0.5.

This model was developed to assist in predicting outcomes for patients who had already undergone prophylactic ileostomy, using clinical and pathological data available shortly after surgery. Therefore, it is not intended for preoperative use (e.g., deciding whether to defunction), but rather for postoperative decision-making regarding the timing and safety of ileostomy closure.

# **Visualizing data**

"Shapviz" is an R package for interpreting ML model predictions, which provides visual explanations based on SHAP (SHapley Additive explanations) values. The SHAP value explains how much each feature contributes to the model's predictions, and whether it is positive or negative. The feature importance plot was used to display the features with the greatest impact on model predictions. Feature importance is ranked based on the average absolute value of the SHAP value. Two samples were selected to create a forced plot of SHAP values for predicting and interpreting the one-sample model.

# **Statistics analysis**

Categorical data were described as frequency and composition ratio. Continuous data that conform to a normal distribution were presented as mean±standard deviation. Non-normally distributed data were presented as frequency (percentage). The chi-square test was used for comparing categorical data, and the t-test for continuous data. A 2-sided value of P < 0.05 was considered statistically significant. R language was used for organizing, analyzing, and visualizing data.

# Results

## **Patient characteristics**

Of the 442 patients included in the analysis, 305 experienced delayed ileostomy closure (69.0%). There were no significant differences between the 2 groups in sex distribution, distant metastasis, tumor pathological type, anastomotic leakage, LARS, and other postoperative complications (all, P>0.05). There were significant differences between the 2 groups in age, BMI, distance between the tumor and the anal verge, lymph node metastasis, depth of tumor infiltration, neoadjuvant chemoradiotherapy, postoperative chemotherapy, and anastomotic stenosis (all, P<0.05) (Table 1).

# **Feature selection**

A total of 14 variables were included in the analysis: LASSO regression was used for the selection of variables due to the fact that too many variables may lead to overfitting of the model, as well as covariance problems (Fig. 1). By choosing lambda values that were 1 standard deviation from the minimum lambda value (within 1 standard error of the minimum value), we identified the 7 most predictive variables (Fig. 2). After screening out the 7 variables with LASSO regression, we fitted a stepwise multifactor logistic regression based on the variables to screen them again. Ultimately, age was excluded because it was not an independent risk factor. Finally, 6 independent variables showed the strongest associations with the dependent variables, ensuring model simplicity and mitigating overfitting problems. The 6 variables were BMI, tumor distance from the anal verge, depth of tumor infiltration, neoadjuvant radiotherapy, postoperative chemotherapy, and anastomotic stenosis (Table 2). Using multivariate logistic regression, we determined that all of these variables were independent risk factors for delayed ileostomy closure.

# Model performance

The performance evaluation metrics of the model are shown in Table 2. The XGBoost model had high area under the ROC curve (AUC) for predicting delayed ileostomy closure in the training set and the validation set: training set AUC = 0.744, 95% confidence interval (CI):

Features	Total (n = 442)	No delayed closure (n = 137)	Delayed closure (n = 305)	р
Gender, n (%)				0.498
Male	156 (35)	52 (38)	104 (34)	
Female	286 (65)	85 (62)	201 (66)	
Age, n (%)				0.015*
≥18~<40	23 (5)	5 (4)	18 (6)	
≥40~<60	206 (47)	52 (38)	154 (50)	
≥60	213 (48)	80 (58)	133 (44)	
BMI, n (%)				< 0.001***
<24	294 (67)	108 (79)	186 (61)	
≥24	148 (33)	29 (21)	119 (39)	
Tumor distance from the anal verge, n (%)				0.005**
≥5 cm	377 (85)	127 (93)	250 (82)	
<5 cm	65 (15)	10 (7)	55 (18)	
Distant metastasis, n (%)				0.613
No	370 (84)	117 (85)	253 (83)	
Yes	72 (16)	20 (15)	52 (17)	
Lymph node, n (%)				0.021*
No	328 (74)	112 (82)	216 (71)	
Yes	114 (26)	25 (18)	89 (29)	
Depth of tumor infiltration, n (%)				0.002**
-	223 (50)	85 (62)	138 (45)	
III-IV	219 (50)	52 (38)	167 (55)	
Pathological type, n (%)				0.269
Adenocarcinoma	420 (95)	132 (96)	288 (94)	
Mucinous adenocarcinoma	15 (3)	2 (1)	13 (4)	
Signet ring cell carcinoma	7 (2)	3 (2)	4 (1)	
Neoadjuvant chemotherapy, n (%)				< 0.001***
No	112 (25)	58 (42)	54 (18)	
Yes	330 (75)	79 (58)	251 (82)	
Postoperative chemotherapy, n (%)				< 0.001***
No	138 (31)	65 (47)	73 (24)	
Yes	304 (69)	72 (53)	232 (76)	
Anastomotic leakage, n (%)				0.058
Features	Total (n = 442)	No delayed closure ( $n = 137$ )	Delayed closure ( $n = 305$ )	р
No	385 (87)	126 (92)	259 (85)	
Yes	57 (13)	11 (8)	46 (15)	
Anastomotic stenosis, n (%)				0.004**
No	401 (91)	133 (97)	268 (88)	
Yes	41 (9)	4 (3)	37 (12)	
LARS, n (%)				0.2
No	299 (68)	99 (72)	200 (66)	
Yes	143 (32)	38 (28)	105 (34)	
Complications, n (%)	. /	· ·	· ·	1
No	373 (84)	116 (85)	257 (84)	
Yes	69 (16)	21 (15)	48 16)	
	\/	<-/p>	,	

Table 1	Comparison	of basic information	between case grou	p and control group

\*:*p* < 0.05 \*\*: *p* < 0.01 \*\*\*: *p* < 0.001

0.686–0.806; test set AUC = 0.809, 95% CI: 0.728–0.889 (Fig. 3A-B). The model had high sensitivity, but low specificity in the training and validation sets: PPV (training set: 0.753, 95% CI: 0.701–0.805; validation set: 0.782, 95% CI: 0.705–0.859), NPV (training set: 0.698, 95% CI: 0.56–0.835; validation set: 0.773, 95% CI: 0.598–0.948). The F1 score of the training set was 0.836, and of the validation set was 0.856) (Table 3).

We assessed the accuracy of the XGBoost model in predicting the probability of delayed ileostomy closure by analyzing the calibration curves and clinical decision curves of the training and validation sets. The calibration



Fig. 1 LASSO coefficient path diagram

curves for both the training and validation sets showed a good fit, indicating that the model predictions are highly consistent with the actual incidence (Fig. 4A-B). The clinical decision curves of the training and validation sets showed that the model has a good degree of clinical utility (Fig. 5A-B).

# Interpretation of the model

The SHAP summary plot of the model explains the effect of each variable on the model (Fig. 6). The order of importance of each variable is, in descending order, BMI > postoperative chemotherapy>distance of the tumor from the anal verge > depth of tumor infiltration > neoadjuvant radiotherapy>anastomotic stenosis. SHAP force diagrams are commonly used to explain how an XGBoost model assesses patient individualized variable contributions. We used the SHAP force plot to explain the level of variable contribution for 2 patients in this study. The colors indicate the contribution of each variable, with blue indicating that the feature is a negative predictor variable (arrow to the left, SHAP value decreases), and red indicating that the feature is a positive predictor variable (arrow to the right, SHAP value increases). The length of the color bar indicates the strength of the contribution, and E [f (x)] indicates the SHAP reference value, which is the average of the model predictions. For the "true positive" patient group, the XGBoost model predicted a delayed closure SHAP value of 0.837, exceeding the baseline value and indicating the occurrence of delayed closure (Fig. 7A). For the group of patients classified as "true negative", the XGBoost model predicted a delayed closure value of 3.12, which did not exceed the reference value thus indicating that no delayed closure occurred (Fig. 7B).

# Discussion

There is currently no consensus on when a temporary loop ileostomy should be closed. Generally, closure is performed at 8–12 weeks after the operation, depending on the situation [18, 31]. In China, patients with a temporary ileostomy usually undergo examinations such as fiber colonoscopy, pelvic magnetic resonance imaging (MRI), defecography, and anorectal pressure measurement to evaluate tumor activity and perianal function before ileostomy closure. The aforementioned examinations take about 1 week. For this study, we defined delayed ileostomy closure as  $closure \ge 90$  days after the index operation based on prior literature suggesting



Fig. 2 LASSO regularization path diagram, where the model has fewer features. while maintaining predictive performance when the regularization parameter value is set to one standard deviation

Table 2	Multivariate	loaistics re	aression of	model	variables
	manuface	iogistics ic	giession or	1110000	vanabics

		-			
	В	SE	Wald	OR_with_Cl	Р
(Intercept)	-0.818	0.313	6.843	0.441(0.236~0.806)	0.009**
BMI	0.824	0.303	7.387	2.281(1.276~4.212)	0.007**
Tumor distance from the anal	1.137	0.482	5.558	3.118(1.291~8.808)	0.018*
Depth of tumor infiltration	0.555	0.279	3.956	1.741(1.011~3.024)	0.047*
Neoadjuvant chemotherapy	0.716	0.310	5.341	2.047(1.113~3.765)	0.021*
Postoperative chemotherapy	0.620	0.300	4.273	1.859(1.03~3.348)	0.039*
Anastomotic stenosis	1.203	0.584	4.249	3.329(1.165~12.106)	0.039*
*. n < 0.05 **. n <	0 01 *** (	0.001			

\*:*p* < 0.05 \*\*: *p* < 0.01 \*\*\*: *p* < 0.001

that 3 months is a widely accepted threshold for planned closure [32]. To our knowledge, this is the first study to develop a SHAP-interpretable XGBoost model to predict the occurrence of temporary ileostomy closure after rectal cancer surgery.

Training and validation sets were used to evaluate the performance of the XGBoost model, and the AUC of the training set for predicting delayed closure was 0.744 (95% CI: 0.686–0.806) and the AUC of the validation set was 0.809 (95% CI: 0.728–0.889). Thus, the model has the potential to assist in the management of patients with

a temporary ileostomy by predicting the risk of delayed closure. Additionally, in order to achieve visualization of the data and results, we introduced the SHAP method to attempt to interpretate the model constructed in this study. The order of importance of each variable is, in descending order, BMI > postoperative chemotherapy > distance of the tumor from the anal verge > depth of tumor infiltration > neoadjuvant radiotherapy > anastomotic stenosis.

Previous studies have revealed that temporary ileostomy closure timing is influenced by various factors, and that there are risk factors associated with delayed closure. In this study, SHAP identified BMI as the most important variable, followed by postoperative chemotherapy and distance of the tumor from the anal verge as the second and third most important variables. Obesity has always been a risk factor for many diseases, and studies have shown that obesity (BMI  $\ge$  30.0 kg/m<sup>2</sup>) can cause various stoma related complications, thereby affecting the timely closure of the stoma [33–34]. Our results showed that a BMI  $\ge$  24.0 kg/m<sup>2</sup> is a significant risk factor for delayed closure.

Postoperative radiotherapy and chemotherapy are considered a risk factors for delayed ileostomy closure [34– 36]. Compared to postoperative adjuvant chemotherapy, little study has been done to examine the impact of neoadjuvant chemoradiotherapy on the delay of ileostomy closure. den Dulk et al. [35] concluded that neoadjuvant



Fig. 3 Receiver operating characteristic curve of the model on the training set(A). Receiver operating characteristic curve of the model validation set(B)

**Table 3** Evaluation of XGBoost model performance in thetraining set and testing set

	Training set	Testing set
AUC	0.744(0.686-0.802)	0.809(0.728–0.889)
Sensitivity	0.939(0.907-0.971)	0.945(0.898–0.992)
Specificity	0.312(0.220-0.405)	0.415(0.264-0.565)
PPV	0.753(0.701-0.805)	0.782(0.705–0.859)
NPV	0.698(0.560-0.835)	0.773(0.598–0.948)
F1 score	0.836	0.856

AUC: area under the receiver operating characteristic curve. PPV: Positive predictive value. NPV: Negative predictive value

radiochemotherapy significantly delays ileostomy closure, probably because preoperative radiochemotherapy increases the risk of postoperative anastomotic complications in rectal cancer patients, which delays the closure of the ileostomy. In this study, the proportion of patients who underwent neoadjuvant therapy and postoperative adjuvant chemotherapy was significantly higher in the delayed closure group, and the LASSO regression analysis showed that postoperative adjuvant chemotherapy and neoadjuvant radiotherapy were risk factors for delayed ileostomy closure. Ileostomy closure is considered by most patients, families, and physicians to be an elective, non-essential procedure, and therefore is often considered unimportant. As such, during postoperative adjuvant chemotherapy, most patients and treating



Fig. 4 Calibration curve of the model training set(A). Calibration curve of the model validation set (B)



Fig. 5 Clinical decision curve of the model on the training set(A). Clinical decision curve of the model validation set(B)

physicians will agree to delay the timing of ileostomy closure for the sake of continuity of chemotherapy. However, a recent study has shown that delaying stoma closure beyond 6 months is associated with nearly a threefold increase in the risk of reoperation [37]. In addition, postoperative chemotherapy can cause patients to become weak, making it difficult for them to undergo surgery in a short period of time and delaying the return of the stoma.

According to reports, 36% of rectal anastomotic leaks can lead to the formation of anastomotic leakage channels, delaying the reintroduction of prophylactic stomas and even causing them to evolve into permanent stomas [38–39]. However, in our study there was no difference in the occurrence of anastomotic leakage between the 2 groups, which may be due to an insufficient number of patients. However, a factor related to anastomotic leakage, the distance between the tumor and the anal margin, was shown to be an independent factor for delayed closure. Research has shown that large tumors near the anus are predictive for irreversible or delayed reversal of a shunt stoma [40]. This may be due to the tumor being closer to the anal margin, which may result in greater tension at the anastomotic site and poor local healing, leading to decisions to delay closure. It may also be due to the fact that with the improvement of surgical techniques and treatment, anastomotic leaks are easy to recognize early and thus controlled without related complications. It is noteworthy that anastomotic stenosis was found to be an independent risk factor in this study, a finding different than in some prior studies. Rectal anastomotic stenosis is the second most serious complication related to anastomosis after rectal cancer resection, and a recent meta-analysis reported that the incidence of anastomotic stenosis after rectal cancer surgery is 17% (95% CI: 13-21%) [41]. Anastomotic stenosis can also delay ileostomy closure and increase the risk of stoma-related complications [13].

Many prior studies have examined risk factors for delayed ileostomy closure after surgery for rectal cancer, but almost all have been limited by using multiple linear regression for model construction [34, 42–43]. Machine learning is a technique that uses computer algorithms to improve model accuracy and precision, enabling better handling of nonlinear relations between multiple variables [25]. Unlike previous studies, we first used LASSSO regression for variable screening. This feature selection method has good interpretability, and can automatically select important features related to the target variable to avoid overfitting of the model [44]. Then, XGBoost ML was used to construct the model, which can improve the accuracy and generalization ability of models, and handle high-dimensional data and complex classification problems [45]. The SHAP method was used to interpret the model, and visualize the results. Multi-center clinical trial data is needed to further evaluate and clarify the accuracy and precision of this predictive model. The model could be integrated into clinical workflows to stratify patients based on their predicted risk of delayed ileostomy closure. For those identified as high-risk, clinicians might consider early postoperative imaging, more frequent follow-up, or prioritizing them for timely stoma closure when appropriate. Applying the model in this way may help support more proactive patient management and optimize use of clinical resources.

Another important consideration is the risk of stoma permanence associated with defunctioning stomas, which may occur even in patients initially intended for reversal [46]. In addition, the presence of a stoma, particularly high-output ileostomies, has been associated to an increased risk of acute kidney injury. These reveal



Fig. 6 Importance chart for SHAP variables

the need for careful monitoring and early closure where appropriate [47].

# Limitations

This study has various limitations to consider. First, we did not use systematic retrieval to include as many risk variables as possible. The study focused on the limited parameters we are familiar with, without considering more potential influencing factors and subgroups, such as diabetes and hypertension. Second, our XGBoost model was trained and validated using data from a single center without external or temporal validation, so its generalizability is unknown. While the current results demonstrate good internal performance, it remains unclear how well the model would perform across different institutions, patient populations, or care settings. We

acknowledge this limitation and plan to conduct prospective external validation in future studies, ideally across multiple centers with more diverse patient cohorts. Third, the sample size used is relatively small. Although model performance was evaluated through training and validation sets, sample size may limit the generalization ability of the results. Finally, although the use of SHAP improves the interpretability of the model, comprehension of ML models remains a challenge for persons not experienced with ML.

# Conclusion

The XGBoost ML model we built helps predict delayed closure of a temporary ileostomy after rectal cancer surgery. In addition, the SHAP method is able to explain the results generated by the ML model, improving the

# A



Fig. 7 SHAP force plot for "true positive" (A). SHAP force plot for "true negative" (B)

interpretability of the model and helping to provide explanations for the source of the predicted results for each patient. This model may help to avoid delayed ileostomy closure after rectal cancer surgery, and help practitioners better manage a temporary ileostomy.

#### Supplementary Information

The online version contains supplementary material available at https://doi.or g/10.1186/s12957-025-03843-w.

Supplementary Material 1

#### Acknowledgements

We would like to thank the patients, doctors, and nurses who participated in this study.

#### Author contributions

Jianying Liu: Writing– review & editing, Writing– original draft, Methodology, Data curation, Conceptualization. Mengxiao Jiang: Investigation, Methodology, Software, Data curation. Yonglan Ge: Investigation, Methodology, Data curation. Zongxin Zheng, Xia Yang & Xiaoping Chen: Data collection, Data input, and verification. Wenhao Zhou: Research and Design Consulting. Huiting Zhang & Meichun Zheng: Visualization, Validation, Conceptualization. Baojia Luo: Visualization, Validation, Methodology, Formal analysis.

#### Data availability

No datasets were generated or analysed during the current study.

# Declarations

# Ethics approval and consent to participate

This study was approved by the Ethics Review Committee of Sun Yat-sen University Cancer Center (SL-B2024-578-01).

## Competing interests

The authors declare no competing interests.

## **Conflict of interest**

The authors declare that they have no known competing financial interests or personal relationships that might influence the work reported in this paper.

## Sources of funding

This study has not attained any funding.

#### Author details

<sup>1</sup>Department of Colorectal Surgery, State Key Laboratory of Oncology in South China, Guangdong Provincial Clinical Research Center for Cancer, Sun Yat-sen University Cancer Center, Guangzhou 510060, P.R. China <sup>2</sup>Department of Urinary Surgery, State Key Laboratory of Oncology in South China, Guangdong Provincial Clinical Research Center for Cancer, Sun Yat-sen University Cancer Center, Guangzhou 510060, P.R. China <sup>3</sup>Nursing Department of Huangpu Yard, State Key Laboratory of Oncology in South China, Guangdong Provincial Clinical Research Center for Cancer, Sun Yat-sen University Cancer Center, Guangzhou 510060, P.R. China

# Received: 27 February 2025 / Accepted: 2 May 2025 Published online: 14 May 2025

## References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA CANCER J CLIN. 2018.
- Heald RJ, Husband EM, Ryall RDH. The mesorectum in rectal cancer surgery the clue to pelvic recurrence? Br J Surg. 1982;69:613–6.
- You YN, Hardiman KM, Bafford A, Poylin V, Francone TD, Davis K et al. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Rectal Cancer. Dis Colon Rectum [Internet]. 2020;63. Available from: https://journals.lww.com/dcrjournal/fulltext/2020/09000/the \_american\_society\_of\_colon\_and\_rectal\_surgeons.6.aspx
- Kverneng Hultberg D, Svensson J, Jutesten H, Rutegård J, Matthiessen P, Lydrup M-L et al. The Impact of Anastomotic Leakage on Long-term Function

After Anterior Resection for Rectal Cancer. Dis Colon Rectum [Internet]. 2020;63. Available from: https://journals.lww.com/dcrjournal/fulltext/2020/05 000/the\_impact\_of\_anastomotic\_leakage\_on\_long\_term.10.aspx

- Keller DS, Talboom K, Van Helsdingen CPM, Hompes R. Treatment modalities for anastomotic leakage in rectal Cancer surgery. Clin Colon Rectal Surg. 2021;34:431–8.
- CHAND M, NASH GF, TALBOT RW. Timely closure of loop ileostomy following anterior resection for rectal cancer. Eur J Cancer Care (Engl). 2008;17:611–5.
- Blok RD, Stam R, Westerduin E, Borstlap WAA, Hompes R, Bemelman WA, et al. Impact of an institutional change from routine to highly selective diversion of a low anastomosis after TME for rectal cancer. Eur J Surg Oncol. 2018;44:1220–5.
- Pisarska M, Gajewska N, Małczak P, Wysocki M, Witowski J, Torbicz G, et al. Defunctioning ileostomy reduces leakage rate in rectal cancer surgery systematic review and meta-analysis. Oncotarget. 2018;9:20816–25.
- Garg PK, Goel A, Sharma S, Chishi N, Gaur MK. Protective diversion stoma in low anterior resection for rectal cancer: A Meta-Analysis of randomized controlled trials. Visc Med. 2019;35:156–60.
- Luglio G, Pendlimari R, Holubar SD, Cima RR, Nelson H. Loop ileostomy reversal after Colon and rectal surgery: A single institutional 5-Year experience in 944 patients. Arch Surg. 2011;146:1191–6.
- Chapman WC, Subramanian M, Jayarajan S, Makhdoom B, Mutch MG, Hunt S, et al. First, do no harm: rethinking routine diversion in Sphincter-Preserving rectal Cancer resection. J Am Coll Surg. 2019;228:547–e5568.
- Gu W, Wu S. Meta-analysis of defunctioning stoma in low anterior resection with total mesorectal excision for rectal cancer: evidence based on thirteen studies. World J Surg Oncol. 2015;13:9.
- 13. Danielsen AK, Park J, Bock D, Haglind E. Early closure of a temporary ileostomy in patients with rectal Cancer. Ann Surg. 2016;265.
- Ye H, Huang S, Yu J, Zhou Q, Xi C, Cao L, et al. Comparison of the clinical outcomes of skin Bridge loop ileostomy and traditional loop ileostomy in patients with low rectal cancer. Sci Rep. 2021;11:9101.
- Jabbal IS, Spaulding AC, Lemini R, Borkar SR, Stanek K, Colibaseanu DT. Temporary vs. permanent Stoma: factors associated with the development of complications and costs for rectal cancer patients. Int J Colorectal Dis. 2022;37:823–33.
- Mehboob A, Perveen S, Iqbal M, Moula Bux K, Waheed A. Frequency and Complications of Ileostomy. Cureus [Internet]. 2020 [cited 2024 Sep 27]; Available from: https://www.cureus.com/articles/41166-frequency-and-complicati ons-of-ileostomy
- Hughes DL, Cornish J, Morris C, On behalf of the LARRIS Trial Management Group. Functional outcome following rectal surgery—predisposing factors for low anterior resection syndrome. Int J Colorectal Dis. 2017;32:691–7.
- Podda M, Coccolini F, Gerardi C, Castellini G, Wilson MSJ, Sartelli M, et al. Early versus delayed defunctioning ileostomy closure after low anterior resection for rectal cancer: a meta-analysis and trial sequential analysis of safety and functional outcomes. Int J Colorectal Dis. 2022;37:737–56.
- Park J, Danielsen AK, Angenete E, Bock D, Marinez AC, Haglind E, et al. Quality of life in a randomized trial of early closure of temporary ileostomy after rectal resection for cancer (EASY trial). Br J Surg. 2018;105:244–51.
- Shimizu H, Yamaguchi S, Ishii T, Kondo H, Hara K, Takemoto K, et al. Who needs diverting ileostomy following laparoscopic low anterior resection in rectal cancer patients? Analysis of 417 patients in a single Institute. Surg Endosc. 2020;34:839–46.
- van den Hout WB, van den Brink M, Stiggelbout AM, van de Velde CJH, Kievit J. Cost-effectiveness analysis of colorectal cancer treatments. Eur J Cancer. 2002;38:953–63.
- 22. Schwalbe N, Wahl B. Artificial intelligence and the future of global health. Lancet. 2020;395:1579–86.
- Kolker E, Özdemir V, Kolker E. How healthcare can refocus on its Super-Customers (Patients, n = 1) and customers (Doctors and Nurses) by leveraging lessons from Amazon, Uber, and Watson. OMICS J Integr Biol. 2016;20:329–33.
- Lei L, Zhang S, Yang L, Yang C, Liu Z, Xu H, et al. Machine learning-based prediction of delirium 24 h after pediatric intensive care unit admission in critically ill children: A prospective cohort study. Int J Nurs Stud. 2023;146:104565.
- Ali S, Akhlaq F, Imran AS, Kastrati Z, Daudpota SM, Moosa M. The enlightening role of explainable artificial intelligence in medical & healthcare domains: A systematic literature review. Comput Biol Med. 2023.
- Kuo C-Y, Kuo L-J, Lin Y. Artificial intelligence based system for predicting permanent stoma after sphincter saving operations. Sci Rep. 2023;13:16039.

- Back E, Häggström J, Holmgren K, Haapamäki MM, Matthiessen P, Rutegård J, et al. Permanent stoma rates after anterior resection for rectal cancer: risk prediction scoring using preoperative variables. Br J Surg. 2021;108:1388–95.
- Wei W, Li Y, Huang T. Using machine learning methods to study colorectal Cancer tumor Micro-Environment and its biomarkers. Int J Mol Sci. 2023;24:11133.
- 29. Li Y, Wu X, Yang P, Jiang G, Luo Y. Machine learning for lung Cancer diagnosis, treatment, and prognosis. Genomics Proteom Bioinf. 2022;20:850–66.
- Zuo D, Yang L, Jin Y, Qi H, Liu Y, Ren L. Machine learning-based models for the prediction of breast cancer recurrence risk. BMC Med Inf Decis Mak. 2023;23:276.
- 31. Hallböök O, Matthiessen P, Leinsköld T, Nyström P-O, Sjödahl R. Safety of the temporary loop ileostomy. Colorectal Dis. 2002;4:361–4.
- Chow A, Tilney HS, Paraskeva P, Jeyarajah S, Zacharakis E, Purkayastha S. The morbidity surrounding reversal of defunctioning ileostomies: a systematic review of 48 studies including 6,107 cases. Int J Colorectal Dis. 2009;24:711–23.
- Chun LJ, Haigh PI, Tam MS, Abbas MA. Defunctioning Loop Ileostomy for Pelvic Anastomoses: Predictors of Morbidity and Nonclosure. Dis Colon Rectum [Internet]. 2012;55. Available from: https://journals.lww.com/dcrjournal/fullte xt/2012/02000/defunctioning\_loop\_ileostomy\_for\_pelvic.10.aspx
- Sier MF, Van Gelder L, Ubbink DT, Bemelman WA, Oostenbroek RJ. Factors affecting timing of closure and non-reversal of temporary ileostomies. Int J Colorectal Dis. 2015;30:1185–92.
- den Dulk M, Smit M, Peeters KC, Kranenbarg EM-K, Rutten HJ, Wiggers T, et al. A multivariate analysis of limiting factors for stoma reversal in patients with rectal cancer entered into the total mesorectal excision (TME) trial: a retrospective study. Lancet Oncol. 2007;8:297–303.
- Lordan JT, Heywood R, Shirol S, Edwards DP. Following anterior resection for rectal cancer, defunctioning ileostomy closure May be significantly delayed by adjuvant chemotherapy: a retrospective study. Colorectal Dis. 2007;9:420–2.
- Munshi E, Segelman J, Matthiessen P, Park J, Rutegård M, Sjöström O, et al. Increased risk of postoperative complications after delayed stoma reversal: a multicenter retrospective cohort study on patients undergoing anterior resection for rectal cancer. Int J Colorectal Dis. 2025;40:36.
- van Koperen PJ, van der Zaag ES, Omloo JMT, Slors JFM, Bemelman WA. The persisting presacral sinus after anastomotic leakage following anterior resection or restorative proctocolectomy. Colorectal Dis. 2011;13:26–9.
- Waterland P, Goonetilleke K, Naumann DN, Sutcliff M, Soliman F. Defunctioning ileostomy reversal rates and reasons for delayed reversal: does delay impact on complications of ileostomy reversal? A study of 170 defunctioning ileostomies. J Clin Med Res. 2015;7:685–9.
- 40. Fukui R, Nozawa H, Hirata Y, Kawai K, Hata K, Tanaka T, et al. Low preoperative maximum squeezing pressure evaluated by anorectal manometry is a risk factor for non-reversal of diverting stoma. Langenbecks Arch Surg. 2021;406:131–9.
- 41. He F, Yang F, Chen D, Tang C, Woraikat S, Xiong J, et al. Risk factors for anastomotic stenosis after radical resection of rectal cancer: A systematic review and meta-analysis. Asian J Surg. 2024;47:25–34.
- Chau TC-Y, Nguyen H, Robertson IK, Harvey X, Tan B, Tan M, et al. Factors affecting timing of loop ileostomy closure: a regional centre's experience with 106 patients. ANZ J Surg. 2024;94:193–8.
- Kim MJ, Kim YS, Park SC, Sohn DK, Kim DY, Chang HJ, et al. Risk factors for permanent stoma after rectal cancer surgery with temporary ileostomy. Surgery. 2016;159:721–7.
- 44. Xie Y, Shi H, Han B. Bioinformatic analysis of underlying mechanisms of Kawasaki disease via weighted gene correlation network analysis (WGCNA) and the least absolute shrinkage and selection operator method (LASSO) regression model. BMC Pediatr. 2023;23:90.
- 45. Zhang Z, Jung C. GBDT-MO: Gradient-Boosted decision trees for multiple outputs. IEEE Trans Neural Netw Learn Syst. 2021;32:3156–67.
- Holmgren K, Häggström J, Haapamäki MM, Matthiessen P, Rutegård J, Rutegård M. Defunctioning stomas May reduce chances of a stomafree outcome after anterior resection for rectal cancer. Colorectal Dis. 2021;23:2859–69.

 Rutegård M, Häggström J, Back E, Holmgren K, Wixner J, Rutegård J, et al. Defunctioning loop ileostomy in anterior resection for rectal cancer and subsequent renal failure: nationwide population-based study. BJS Open. 2023;7:zrad010.

# **Publisher's note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.